

YEAR 12 – TRIAL 2006 – EXTENSION 1**QUESTION 1****MARKS**

a) Find $\frac{d}{dx}(e^{2x} \cos 3x)$ 2

b) Evaluate 2

$$\int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\sqrt{1-3x^2}} dx$$

c) Evaluate $\lim_{x \rightarrow 0} \frac{x}{\sin 4x}$ 1

d) Use the substitution $u = 1 + x^4$ to evaluate 2

$$\int_0^1 \frac{x^3}{1+x^4} dx$$

e) If the roots of the equation $x^3 + 2x^2 - x - p = 0$ are c , $c + 1$ and $c + 3$, find the value of p . 2

f) A (1, 0) and B (2, 4) are two points in the number plane. 1
Find the coordinates of the point P that divides the interval AB externally in the ratio 3:1.

g) Find the term independent of x in the expansion $\left(x + \frac{2}{x}\right)^{10}$ 2

QUESTION 2**MARKS**

a) Solve $\frac{3}{2-x} \geq 1$ 2

b) Consider the polynomial 2

$$P(x) = (x+2)(x-3)Q(x) + ax + b$$

Given that $P(x)$ has remainders 1 and 6 when divided by $(x+2)$ and $(x-3)$ respectively, find a and b .

c) Consider the function

$$y = \pi + 2 \sin^{-1}\left(\frac{x}{3}\right)$$

i) Find the domain and the range. 2

ii) Sketch the graph of the function. 2

d) The region bounded by the curve $y = \sin^{-1} x$, the y axis and $y = \frac{\pi}{6}$, is rotated about the y axis to form a solid.


i) Show that the volume of the solid obtained is given by 2

$$V = \pi \int_0^{\frac{\pi}{6}} \sin^2 y \, dy$$

ii) Find the volume of the solid. 2

QUESTION 3**MARKS**

- a) A, X, B and Y are points on the circumference of a circle such that $AX = AY$.
 XY meets AB at the point M.
 The tangent at B meets the chord XY produced at T.



- | | |
|--|---|
| i) Explain why $\angle ABY = \angle AYX$. | 1 |
| ii) Show that AB bisects $\angle XBY$. | 1 |
| iii) Show that $BT = MT$. | 2 |
|
b) Consider the functions $f(x) = 4 - x^2$ and $g(x) = \ln x$. | |
| i) Sketch the graphs of $f(x)$ and $g(x)$ on the same set of axes for $x > 0$. | 2 |
| ii) Use your graph to show that the equation $\ln x + x^2 - 4 = 0$ has only one root which is near $x = 1.5$ | 1 |
| iii) Use one application of Newton's method to find a better approximation of the root of the equation $\ln x + x^2 - 4 = 0$ | 2 |
|
c) Ten people are to be seated around two circular tables. Six of them can sit at one table and the remaining four can sit at the other table. | |
| i) How many different groups can be formed to sit around these two tables? | 1 |
| ii) How many seating arrangements are possible around the two tables? | 2 |

QUESTION 4**MARKS**

- a) The population N of a particular species of birds in an island at any time t is expressed as

$$N = 2000 + ce^{-kt}, \text{ where } c \text{ and } k \text{ are constants.}$$

Given that the initial population was 11 000 birds and it decreased to 8 000 after 10 years,

- i) find the constants c and k . 2
- ii) find the time required for the population to decrease to 6 000 birds. 2

- b) Two points $P(2ap, ap^2)$ and $Q(2aq, aq^2)$ lie on the parabola $x^2 = 4ay$. The tangents to the parabola at P and Q intersect at an angle of 60° at a point A .

- i) Show that the coordinates of A are $(a(p+q), apq)$ 2

(You may assume that the equation of the tangent at P is $y = px - ap^2$)

- ii) Show that $p - q = \sqrt{3}(1 + pq)$ 1

- iii) Show that as P and Q moves on the parabola the point A moves on the curve with equation $x^2 = 3a^2 + 10ay + 3y^2$ 2

- c) 75% of the workers in a shoe factory are highly skilled, while 25% are less skilled.

Each worker makes the same number of pairs of shoes a day.

For highly skilled workers only 1% of the pairs of shoes are defective, and 2% are defective for the less skilled.

- i) What is the probability that a pair of shoes made in this factory is defective? 1
- ii) What is the probability that, in a random parcel of 20 pairs of shoes, no more than one pair is defective? 2

QUESTION 5**MARKS**

- a) Let $g(x) = \frac{x}{1+x^2}$ for all real values of x .
- i) Sketch the graph of $g(x)$ showing the coordinates of the turning points and the x and y intercepts. 3
- ii) What is the largest domain containing the value $x = 2$ for which $g(x)$ has an inverse function $g^{-1}(x)$? 1
- iii) Sketch the graph of the inverse function $y = g^{-1}(x)$ on the same set of axes as your graph in part (i). 1
- iv) Find an expression for $y = g^{-1}(x)$ in terms of x . 1
- b) A particle moves in a straight line and its displacement x centimetres from the origin at time t seconds is given by :
- $$x = 2 + 4 \cos^2 \left(\frac{\pi}{2} t \right)$$
- i) Show that the motion of the particle is simple harmonic. 2
- ii) Find the maximum velocity of the particle. 1
- c) Use mathematical induction to prove that, for all integers n with $n \geq 1$. 3

$$\sum_{k=1}^n k(k+1) = \frac{n(n+1)(n+2)}{3}$$

QUESTION 6**MARKS**

- a) Consider the function $f(x) = \cos^{-1}\left(\frac{1-x}{1+x}\right) - 2 \tan^{-1} \sqrt{x}$ where $x > 0$.


i) Show that $f'(x) = 0$

3

ii) Sketch the graph of $y = f(x)$

1

- b) A tank has the shape of an inverted right circular cone of base radius 8m and height 12m contains water to a depth 3 m



Water starts to be poured into the tank at the constant rate of $1.2 \text{ m}^3/\text{min}$ but at the same time a leak at the vertex of the tank causes a discharge of the water at a constant rate of $0.2 \text{ m}^3/\text{min}$.

- i) Show that the volume V of the water in cubic metres at any time t can be expressed as

2

$$V = 4\pi + t$$

- ii) Calculate the rate at which the height is increasing at the time $t = 28\pi$ minutes.

3

- iii) Calculate the rate at which the area of the water in contact with the wall of this tank is increasing at the time $t = 104\pi$ minutes.

3


QUESTION 7**MARKS**

- a) Given that $(1 + \frac{1}{x})^n (1 + x)^n = \frac{1}{x^n} (1 + x)^{2n}$, show that

3

$${}^nC_0 {}^nC_3 + {}^nC_1 {}^nC_4 + \dots + {}^nC_{n-3} {}^nC_n = \frac{(2n)!}{(n+3)!(n-3)!}$$

b)



Two projectiles are projected at the same time towards a target at point B(d,d). One of the projectiles is projected from O(0,0) with a velocity V, while the second projectile is projected from A(2d,0) with velocity U.

At the instant when the projectiles are projected target B starts falling vertically down under gravity.

Consider the axes as shown and assume that there is no air resistance, and that g is the acceleration due to gravity.

- i) Show that the equations of the positions of the target are

1

$$y = -\frac{1}{2}gt^2 + d \text{ and } x = d$$

- ii) Find the equations of the positions of the two projectiles.

3

- iii) The projectile from O hits the target in the air.

3

Find the time taken by the projectile to hit the target and show that $V > \sqrt{gd}$

- iv) After the impact a fragment of the target falls vertically down under gravity and t minutes later the particle from A hits this fragment.

2

$$\text{Show that } t = \sqrt{2d} \left(\frac{1}{U} - \frac{1}{V} \right)$$

-1-

Question 1:

a) $\frac{d}{dx} (e^{2x} \cos 3x)$

Using the product rule:

$u = e^{2x} \quad v = \cos 3x$

$u' = 2e^{2x} \quad v' = -3 \sin 3x$

$\therefore = 2e^{2x} \cdot \cos 3x - 3e^{2x} \sin 3x$

$= e^{2x} (2 \cos 3x - 3 \sin 3x)$ (2 marks)

b) $\int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\sqrt{1-3x^2}} dx$

$= \int_{-\frac{1}{2}}^{\frac{1}{2}} \frac{1}{\sqrt{3} \sqrt{\frac{1}{3} - x^2}} dx$

$= \frac{1}{\sqrt{3}} \left[\sin^{-1} \left(\frac{x}{\frac{1}{\sqrt{3}}} \right) \right]_{-\frac{1}{2}}^{\frac{1}{2}} = \frac{1}{\sqrt{3}} \left[\sin^{-1} \sqrt{3} x \right]_{-\frac{1}{2}}^{\frac{1}{2}}$

$= \frac{1}{\sqrt{3}} \left[\sin^{-1} \left(\frac{\sqrt{3}}{2} \right) - \sin^{-1} \left(-\frac{\sqrt{3}}{2} \right) \right]$

$= \frac{1}{\sqrt{3}} \left(\frac{\pi}{3} - -\frac{\pi}{3} \right) = \frac{2\pi}{3\sqrt{3}}$ (2 marks)

c) $\lim_{x \rightarrow 0} \frac{x}{\sin 4x} = \lim_{x \rightarrow 0} \frac{1}{4} \times \frac{4x}{\sin 4x}$

$= \lim_{x \rightarrow 0} \frac{1}{4} \times \frac{1}{\frac{\sin 4x}{4x}} = \frac{1}{4} \times \frac{1}{1} = \frac{1}{4}$ (1 mark)

d) $\int_0^1 \frac{x^3}{1+x^4} dx$

let $u = 1+x^4$

$\therefore \frac{du}{dx} = 4x^3$

$\therefore \frac{1}{4} du = x^3 dx$

when $x=1$, $u=2$

when $x=0$, $u=1$

$\therefore I = \frac{1}{4} \int_1^2 \frac{1}{u} du = \frac{1}{4} [\ln u]_1^2$

$= \frac{1}{4} (\ln 2 - \ln 1) = \frac{1}{4} \ln 2$ (2 marks)

e) $x^3 + 2x^2 - x - p = 0$

The roots are $c, c+1, c+3$.

The sum of the roots is $c+c+1+c+3 = -2$

$\therefore 3c+4 = -2, 3c = -6, c = -2$

The products of the roots is

$c(c+1)(c+3) = p$, but $c = -2$

$\therefore -2(-1)(1) = p \therefore p = 2$ (2 marks)

f) $A(1,0) \quad B(2,4)$

$(1,0) \quad -3 \quad x = \frac{(1 \times 1) + (-3 \times 2)}{-3+1}$

$(2,4) \quad 1 \quad = \frac{1-6}{-2} = 2\frac{1}{2}$

$y = \frac{(1 \times 0) + (-3 \times 4)}{-3+1} = \frac{-12}{-2} = 6$

$\therefore$ The point P is $(2\frac{1}{2}, 6)$ (1 mark)

g) $(x + \frac{2}{x})^{10}$

$T_{r+1} = {}^{10}C_r x^{10-r} \left(\frac{2}{x}\right)^r$

$= {}^{10}C_r x^{10-r} 2^r x^{-r}$

$= {}^{10}C_r 2^r x^{10-2r}$

To find the term independent of x , we

let the power of x equal zero.

$\therefore 10-2r = 0, r = 5$

$\therefore T_6 = {}^{10}C_5 2^5 = 8064$ (2 marks)

Question 2:

a) $\frac{3}{2-x} > 1$

$\frac{3}{2-x} - 1 > 0, \frac{3-(2-x)}{2-x} > 0$

$\frac{3-2+x}{2-x} > 0, \frac{1+x}{2-x} > 0$

-2-

x	-1	2
x	-	+
$-x$	+	-
$1/x$	-	+

$\therefore$ solutions are $-1 \leq x \leq 2$. (2 marks)

i) $P(x) = (x+2)(x-3)Q(x) + ax+b$

is $P(-2)=1$ and $P(3)=6$, by

substitution we get $-2a+b=1$

$2a-b=-1$ (1)

$3a+b=6$ (2)

By solving these equations

simultaneously we get $5a=5$

$a=1$. $3+b=6$, $\therefore b=3$

(2 marks)

i) $y = \pi + 2\sin^{-1}\left(\frac{x}{3}\right)$

Domain: $-1 \leq \frac{x}{3} \leq 1$


$-3 \leq x \leq 3$.

range: $-\frac{\pi}{2} \leq \sin^{-1}\left(\frac{x}{3}\right) \leq \frac{\pi}{2}$

$-\pi \leq 2\sin^{-1}\left(\frac{x}{3}\right) \leq \pi$


$\therefore 0 \leq y \leq 2\pi$.

(2 marks)



(2 marks)

d)



i) $y = \sin^{-1}x \therefore x = \sin y \therefore x^2 = \sin^2 y$

Volume formed when the area is

rotated about the y axis is

$V = \pi \int_a^b x^2 dy \therefore V = \pi \int_0^{\pi/2} \sin^2 y dy$ (2 marks)

ii) $\cos 2y = 1 - 2\sin^2 y$

$2\sin^2 y = 1 - \cos 2y$

$\sin^2 y = \frac{1}{2}(1 - \cos 2y)$

$\therefore V = \frac{\pi}{2} \int_0^{\pi/2} (1 - \cos 2y) dy = \frac{\pi}{2} \left[y - \frac{1}{2} \sin 2y \right]_0^{\pi/2}$
 $= \frac{\pi}{2} \left[\frac{\pi}{2} - \frac{\sqrt{3}}{4} - (0 - 0) \right] = \frac{\pi^2}{4} - \frac{\sqrt{3}\pi}{8}$ units³ (2 marks)

Question 3:


a) Data: $AX = AY$

Aim: (i) Explain why $\angle ABY = \angle AXY$

(ii) Show that $AB \perp XY$.

(iii) Show that $BT = MT$.

Construction:



Proof:

i) Let $\angle ABY = \alpha \therefore \angle AXY = \alpha$ (angles at the circumference subtended by the same arc AY are equal).

$AX = AY$ (given)

$\therefore \triangle AXY$ is isosceles (2 equal sides)

$\therefore \angle AXY = \angle AYX = \alpha$ (base angles in isosceles triangle AXY equal)

$\therefore \angle AXY = \angle AYX = \alpha$ (1 mark)

i) $\angle AXY = \angle ABX = \alpha$ (angles at the circumference subtended by same arc AX equal)

$\therefore \angle AXY = \angle ABX = \alpha$

Hence, AB bisects $\angle XBY$. (1 mark)

ii) Let $\angle TBY = \beta$

$\therefore \angle BAY = \beta$ (angle in the alternate

segment, tangent TB and chord BY)

$\therefore \angle BMT = \alpha + \beta$ (exterior angle of

$\triangle AMY$ equals the sum of the opposite interior angles)


$\angle MBT = \angle MBY + \angle YBT = \alpha + \beta$ (adj. $\angle$'s)

$\therefore \angle BMT = \angle MBT = \alpha + \beta$

$\therefore \triangle BMT$ is isosceles (base angles =)

$\therefore BT = MT$ (2 equal sides in isosceles $\triangle BMT$) (2 marks)

i) $f(x) = 4 - x^2$ $g(x) = \ln x$



ii) From the graph, $y = \ln x$ and $y = 4 - x^2$ intersect only once near $x = 1.5$.

The equation $\ln x = 4 - x^2$ (i.e. $\ln x + x^2 - 4 = 0$) has a root close to $x = 1.5$. (1 mark)

iii) Let $h(x) = \ln x + x^2 - 4$

$\therefore h'(x) = \frac{1}{x} + 2x$

$\therefore h(1.5) = \ln(1.5) + (1.5)^2 - 4 = -1.3445 \dots$

$\therefore h'(1.5) = \frac{1}{1.5} + (2 \times 1.5) = 3.666 \dots = 3\frac{2}{3}$

$\therefore x_1 = 1.5 - \frac{h(1.5)}{h'(1.5)} = 1.866 \dots \approx 1.87$ (2 dp)

$\therefore h(1.87) = 0.1228 \dots$

$\therefore x = 1.87$ is a better approximation to the root. (2 marks)

c) There are $^{10}C_6$ ways of selecting 6 people for the first table, leaving 4C_4 ways of selecting 4 people from the remaining 4, for the second table.

$\therefore ^{10}C_6 \times ^4C_4 = 210$ different groups can be selected. (1 mark)

ii) For each group in part (i) there are $5!$ arrangements for one table and $3!$ arrangements for the other.

$\therefore$ Total is $5! \times 3! \times 210 = 151200$ (2 marks)

Question 4:

a) $N = 2000 + ce^{-kt}$

When $t = 0$, $N = 11000$

$\therefore 11000 = 2000 + ce^0$

$\therefore c = 9000$

$\therefore N = 2000 + 9000e^{-kt}$

When $t = 10$, $N = 8000$

$\therefore 8000 = 2000 + 9000e^{-10k}$

$$100 = 9000e^{-10k}$$

$$= e^{-10k}$$

plying $\log_e$ to both sides

$$\ln\left(\frac{2}{3}\right) = -10k \ln e$$

$$k = \frac{-1}{10} \ln\left(\frac{2}{3}\right)$$

$$N = 2000 + 9000e^{\frac{1}{10} \ln\left(\frac{2}{3}\right)t} \quad (2 \text{ marks})$$

$$N = 6000$$

$$6000 = 2000 + 9000e^{\frac{1}{10} \ln\left(\frac{2}{3}\right)t}$$

$$4000 = 9000e^{\frac{1}{10} \ln\left(\frac{2}{3}\right)t}$$

$$= e^{\frac{1}{10} \ln\left(\frac{2}{3}\right)t}$$

plying $\log_e$ on both sides

$$\ln\left(\frac{4}{9}\right) = \frac{1}{10} \ln\left(\frac{2}{3}\right)t \ln e$$

$$t = \frac{\ln\left(\frac{4}{9}\right)}{\frac{1}{10} \ln\left(\frac{2}{3}\right)} = 20 \text{ years} \quad (2 \text{ marks})$$

The equation of the tangent at

$$y = px - ap^2 \text{ (given)}$$

therefore, the equation of

the tangent at Q is $y = qx - aq^2$

$$\text{subtraction } (p-q)x - a(p^2 - q^2) = 0$$

$$(p-q)x = a(p-q)(p+q)$$

$$= a(p+q) \quad (p \neq q)$$

$$y = p[a(p+q)] - ap^2$$

$$= ap^2 + apq - ap^2$$

$$= apq$$

$$A[a(p+q), apq] \quad (2 \text{ marks})$$

The gradient at P is p. The

gradient at Q is q. The lines

intersect at an angle of 60°

$$\tan 60^\circ = \frac{m_1 - m_2}{1 + m_1 m_2}$$

$$\tan 60 = \frac{p-q}{1+pq} \therefore \sqrt{3} = \frac{p-q}{1+pq}$$

$$\text{so } p-q = \sqrt{3}(1+pq) \quad (1 \text{ mark})$$

$$\text{iii) } x = a(p+q)$$

$$\therefore \frac{x}{a} = p+q$$

$$\therefore \left(\frac{x}{a}\right)^2 = (p+q)^2 = p^2 + 2pq + q^2$$

$$= p^2 - 2pq + q^2 + 4pq$$

$$= (p-q)^2 + 4pq$$

$$= [\sqrt{3}(1+pq)]^2 + 4pq$$

$$= 3(1+2pq+p^2q^2) + 4pq$$

$$= 3 + 6pq + 3p^2q^2 + 4pq$$


$$\therefore \frac{x^2}{a^2} = 3 + 10pq + 3p^2q^2$$

$$\text{but } y = apq, \therefore pq = \frac{y}{a}$$

$$\therefore \frac{x^2}{a^2} = 3 + \frac{10y}{a} + \frac{3y^2}{a^2}$$

$$\therefore x^2 = 3a^2 + 10ay + 3y^2 \quad (2 \text{ marks})$$

c) i)



$$\therefore \text{probability (defective)} = (0.75 \times 0.01)$$

$$+ (0.25 \times 0.02) = 0.0125 \quad (1 \text{ mark})$$

$$\text{ii) Probability (defective)} = 0.0125$$

$$\text{Probability (non-defective)} = 1 - 0.0125$$

$$= 0.9875$$

By using binomial probability;

No more than one pair defective

is none defective or 1 defective.

$$\therefore P = {}^{20}C_0 (0.9875)^{20} + {}^{20}C_1 (0.0125 \times 0.9875^{19})$$

$$= 1 \times 0.7775 + (20 \times 0.0125 \times 0.7874 \dots)$$

$$= 0.9744 \dots \quad (2 \text{ marks})$$

Question 5:

1) $g(x) = \frac{x}{1+x^2}$

i) domain: all real x

asymptotic behaviour: $x \rightarrow \pm \infty$

$y \approx \frac{1}{x} \rightarrow 0$



$\therefore y=0$ is a horizontal asymptote.

derivative: $\frac{dy}{dx} = \frac{1+x^2-2x^2}{(1+x^2)^2}$

$\frac{dy}{dx} = \frac{1-x^2}{(1+x^2)^2}$

Let $\frac{dy}{dx} = 0$ to find possible stationary turning points.

$1-x^2=0 \therefore x^2=1 \therefore x=\pm 1$



2) $g(x)$ has an inverse function when it is a one-one function.

That is, for every x -value there

one y -value and for every

y -value there is one x -value.

The largest domain containing $x=2$ for which this occurs is $x \geq 1$. (1 mark)

iii) On graph. (1 mark)

iv) Domain of function: $x \geq 1$

Range of function: $0 < y \leq \frac{1}{2}$

Domain of inverse function: $0 < x < \frac{1}{2}$

Range of inverse function: $y \geq 1$

By interchanging x and y values, we get;

$x = \frac{y}{1+y^2}$, $x(1+y^2)=y$, $y=x+xy^2$

$xy^2 - y + x = 0$ using the quadratic formula,

$y = \frac{1 \pm \sqrt{1-4x^2}}{2x}$

Since the range is $y \geq 1$, y is only

possible when $y = \frac{1 + \sqrt{1-4x^2}}{2x}$ (1 mark)

b) $x = 2 + 4\cos^2\left(\frac{\pi}{2}t\right)$

but $\cos \pi t = 2\cos^2 \frac{\pi}{2}t - 1$

$\therefore 2\cos \pi t = 4\cos^2 \frac{\pi}{2}t - 2$

and $2 + 2\cos \pi t = 4\cos^2 \frac{\pi}{2}t$

$\therefore x = 2 + (2 + 2\cos \pi t)$

$= 4 + 2\cos \pi t$ ①

so $\dot{x} = -2\pi \sin \pi t$ ②

$\ddot{x} = -2\pi^2 \cos \pi t$ ③

$= -\pi^2 (2\cos \pi t)$

From ①, $2\cos \pi t = x - 4$

$\therefore \ddot{x} = -\pi^2 (x - 4)$ period = 2

since the acceleration is proportional to the displacement, where the constant of its proportion is negative.

Hence, the motion is simple harmonic

about $x=4$. (2 marks)

ii) $\dot{x} = -2\pi \sin \pi t$. As maximum value

$\sin \pi t = 1$ then max velocity
 $2\pi \text{ cm/sec.}$ (1 mark)

$$\sum_{k=1}^n k(k+1) = \frac{n(n+1)(n+2)}{3}$$

$$1 \times 2 + 2 \times 3 + 3 \times 4 + \dots + n(n+1) = \frac{n(n+1)(n+2)}{3}$$

p 1: For $n=1$,
 $S = 1 \times 2 = 2$

$$HS = \frac{1(2)(3)}{3} = \frac{6}{3} = 2$$

HS = 2HS $\therefore$ Statement true for $n=1$

p 2: Assuming the statement
 is true for $n=k$.

$$1 \times 2 + 2 \times 3 + 3 \times 4 + \dots + k(k+1) = \frac{k(k+1)(k+2)}{3}$$

aim is to prove the statement

is true for $n=k+1$.

$$1 \times 2 + 3 \times 4 + \dots + k(k+1) + (k+1)(k+2)$$

$$+ (k+1)(k+2)(k+3)$$

$$= 1 \times 2 + 2 \times 3 + \dots + k(k+1) + (k+1)(k+2)$$

$$+ \frac{k(k+1)(k+2)}{3} + \frac{(k+1)(k+2)}{3}$$

$$\frac{(k+1)(k+2)}{3} \left(\frac{k}{3} + 1 \right)$$

$$\frac{(k+1)(k+2)}{3} \left(\frac{k+3}{3} \right)$$

$$\frac{(k+1)(k+2)(k+3)}{3} = k+HS$$

ie, if the statement is true

$n=k$, it is also true for

$(k+1)$.

3: From step 1, statement

is true for $n=1$. Hence by step 2

statement must be true for $n=2$,

hence true for $n=3$, and

is true for all positive integers.

(3 marks)

Question 6:

a) $f(x) = \cos^{-1} \left(\frac{1-x}{1+x} \right) - 2 \tan^{-1} \sqrt{x}$

i) Let $u = \frac{1-x}{1+x} \therefore y = \cos^{-1} u$

$$\frac{dy}{dx} = \frac{-1}{(1+x)^2} \cdot \frac{-1}{(1+x)^2} = \frac{-2}{(1+x)^2}$$

$$\frac{dy}{du} = \frac{-1}{\sqrt{1-u^2}} = \frac{-1}{\sqrt{1-\frac{(1-x)^2}{(1+x)^2}}} = \frac{-1}{\sqrt{\frac{(1+x)^2 - (1-x)^2}{(1+x)^2}}}$$

$$= \frac{-1}{\sqrt{\frac{4x}{(1+x)^2}}} = \frac{-1}{\frac{\sqrt{4x}}{1+x}} = \frac{-1}{\frac{2\sqrt{x}}{1+x}} = \frac{-(1+x)}{2\sqrt{x}}$$

$$\therefore \frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx} = \frac{-(1+x)}{2\sqrt{x}} \times \frac{-2}{(1+x)^2}$$

$$= \frac{1}{\sqrt{x}(1+x)}$$

if $z = 2 \tan^{-1} \sqrt{x}$, let $u = \sqrt{x} \therefore z = 2 \tan^{-1} u$

$$\frac{dz}{dx} = \frac{1}{2} \times \frac{1}{\sqrt{x}} = \frac{1}{2\sqrt{x}}$$

$$\frac{dz}{du} = 2 \times \frac{1}{1+u^2} = \frac{2}{1+x}$$

$$\therefore \frac{dz}{dx} = \frac{dz}{du} \times \frac{du}{dx} = \frac{2}{1+x} \times \frac{1}{2\sqrt{x}} = \frac{1}{\sqrt{x}(1+x)}$$

$$\therefore f'(x) = \frac{1}{\sqrt{x}(1+x)} - \frac{1}{\sqrt{x}(1+x)}$$

$$\therefore f'(x) = 0 \quad (3 \text{ marks})$$


ii) As $f(x)$ is a continuous function in the domain $x > 0$; and $f'(x) = 0$.


$$\therefore f(x) = C. \text{ when } x=1, f(x) = \cos^{-1} 0 - 2 \tan^{-1} 1$$

$$\text{so } f(x) = \frac{\pi}{2} - 2 \times \frac{\pi}{4} = 0$$

$$\therefore f(x) = \frac{\pi}{2} - 2 \times \frac{\pi}{4} = 0$$

$$\therefore f(x) = 0 \text{ for } x > 0$$





Let the water in the tank at a time t be in the shape of a cone of radius r and

but $s = \sqrt{h^2 + r^2}$ (Pythagoras' theorem)
 $= \sqrt{h^2 + \left(\frac{2h}{3}\right)^2}$
 $= \sqrt{h^2 + \frac{4h^2}{9}}$
 $= \sqrt{\frac{13h^2}{9}} = \frac{h}{3} \sqrt{13} \quad (h > 0)$

$A = \pi r s = \pi \times \frac{2h}{3} \times \frac{\sqrt{13}}{3} h = \frac{2\sqrt{13}}{9} \pi h^2$

$\frac{dA}{dh} = \frac{4\sqrt{13}}{9} \pi h$

$\frac{dA}{dt} = \frac{dA}{dh} \cdot \frac{dh}{dt} = \frac{4\sqrt{13}}{9} \pi h \times \frac{9}{4\pi h^2} = \frac{\sqrt{13}}{h}$

$\triangle ABC$ is similar to $\triangle ADE$

(equiangular). Volume of a cone

at a time t is $V = \frac{1}{3} \pi r^2 h$.

By using the ratio of sides of

similar $\triangle$'s $\frac{r}{8} = \frac{h}{12}$, $r = \frac{8h}{12}$

$\therefore r = \frac{2h}{3}$

$\therefore V = \frac{1}{3} \pi h \left(\frac{4h^2}{9}\right)$, $V = \frac{4\pi}{27} h^3$

at $t=0$, $h=3$ m $\therefore V = 4\pi$ m³.

As water enters the tank at a

rate of 1.2 m³/min, and leaks

from the tank at a rate

of 0.2 m³/min, then $\frac{dV}{dt} = 1.2 - 0.2$

$= 1$ m³/min $\therefore V = \int 1 dt = t + C$

But when $t=0$, $V=4\pi$

$\therefore V = (t + 4\pi)$ m³/min. (2 marks)

ii) $\frac{dV}{dt} = \frac{dV}{dh} \cdot \frac{dh}{dt}$, but $V = \frac{4\pi}{27} h^3$ (Part i)

$\therefore \frac{dV}{dh} = \frac{4\pi}{27} \times 3h^2 = \frac{4\pi}{9} h^2$

$\frac{dV}{dt} = 1$

$\therefore 1 = \frac{4\pi}{9} h^2 \cdot \frac{dh}{dt} \therefore \frac{dh}{dt} = \frac{9}{4\pi h^2}$

Now $t=28\pi$, $V=28\pi+4\pi=32\pi$

$\pi h^3 = 32\pi$, $h^3 = 32$, $\therefore h=2$

$\therefore \frac{dh}{dt} = \frac{9}{4\pi \times 2^2} \therefore \frac{dh}{dt} = \frac{9}{16\pi}$ m/min (3 marks)

ii) Area of water in contact with the wall is $A = \pi r s$ (s is slant height)

When $t=12\pi \therefore V=104\pi+4\pi=108\pi$

$\therefore 108\pi = \frac{4\pi}{27} h^3$, $h^3 = 729$, $h=9$ m

$\therefore \frac{dA}{dt} = \frac{\sqrt{13}}{9}$ m²/min (3 marks)

Question 7:

a) LHS = $\left(1 + \frac{1}{x}\right)^n (1+x)^n = (1+x^{-1})^n (1+x)^n$
 $= \left({}^n C_0 + {}^n C_1 x^{-1} + \dots + {}^n C_{n-3} x^{-3} + {}^n C_{n-2} x^{-2} + {}^n C_{n-1} x^{-1} + {}^n C_n\right)$
 $\left({}^n C_0 + {}^n C_1 x + {}^n C_2 x^2 + {}^n C_3 x^3 + {}^n C_4 x^4 + \dots + {}^n C_n x^n\right)$

${}^n C_0 {}^n C_3 + {}^n C_1 {}^n C_4 + \dots + {}^n C_{n-3} {}^n C_n$ would represent

the co-efficient of x^3 in the expansion of LHS.

To get x^3 in RHS we need to get the

coefficient of x^{n+3} in $(1+x)^{2n}$. Trm in

$(1+x)^{2n}$ is ${}^{2n} C_r x^r \therefore$ (coef. of x^{n+3} is obtained

for $r=n+3 \therefore$ coefficient required

is ${}^{2n} C_{n+3}$, but ${}^{2n} C_{n+3} = \frac{(2n)!}{(n+3)!(2n-(n+3))!}$
 $= \frac{2n!}{(n+3)!(n-3)!}$

Hence, by equating the coefficient of

x^3 on each side of the identity.

$\therefore {}^n C_0 {}^n C_3 + {}^n C_1 {}^n C_4 + \dots + {}^n C_{n-3} {}^n C_n = \frac{(2n)!}{(n+3)!(n-3)!}$ (3 marks)

b) For the target at B, there are two

motions: horizontal motion and

vertical motion.

Horizontal motion:

$\ddot{x} = 0$, by integrating with respect

time t , $\dot{x} = C_1$. When $t=0$, $\dot{x}=0$

$$C_1 = 0$$

$$\dot{x} = 0$$

By integrating with respect of

time t , $x = C_2$. When $t=0$, $x=d$

$$x = d$$

Vertical motion:

$\ddot{y} = -g$. By integrating with

respect of time t , $\dot{y} = -gt + K_1$

When $t=0$, $\dot{y}=0 \therefore K_1 = 0$

$$\dot{y} = -gt$$

By integrating with respect

of time t : $y = -\frac{1}{2}gt^2 + K_2$

When $t=0$, $y=d \therefore K_2 = d$

$$y = -\frac{1}{2}gt^2 + d \quad (1 \text{ mark})$$

The initial speed of V is

directed towards (d, d) . Let

α be the angle of v with

positive x -axis.

$$\tan \alpha = \frac{d}{d} = 1 \therefore \alpha = 45^\circ$$

Similarly, the angle of U with

negative x axis is 45° .

The angle of U with positive

x axis is $180^\circ - 45^\circ = 135^\circ$.

Horizontal motion of the projectile

from O .

$\ddot{x} = 0$. By integrating with respect

time t , $\dot{x} = C_1$. When $t=0$,

$$\dot{x} = V \cos 45^\circ = \frac{V}{\sqrt{2}} \therefore C_1 = \frac{V}{\sqrt{2}} \therefore \dot{x} = \frac{V}{\sqrt{2}}$$

By integrating with respect of time t ,

$$x = \frac{V}{\sqrt{2}}t + C_2 \text{ . When } t=0, x=0$$

$$\therefore C_2 = 0 \therefore x = \frac{V}{\sqrt{2}}t + C_2 \text{ . When } t=0, x=0$$

$$\therefore C_2 = 0 \therefore x = \frac{V}{\sqrt{2}}t$$

Vertical motion of the projectile from O .

$\ddot{y} = -g$. By integrating with respect of

time t , $\dot{y} = -gt + K_1$. When $t=0$,

$$\dot{y} = V \sin 45^\circ = \frac{V}{\sqrt{2}} \therefore K_1 = \frac{V}{\sqrt{2}}$$

$$\therefore \dot{y} = -gt + \frac{V}{\sqrt{2}}$$

By integrating with respect of time t .

$$y = -\frac{1}{2}gt^2 + \frac{V}{\sqrt{2}}t + K_2 \text{ . When } t=0, y=0.$$

$$\therefore K_2 = 0 \therefore y = -\frac{1}{2}gt^2 + \frac{V}{\sqrt{2}}t$$

Horizontal motion of the projectile from A .

$\ddot{x} = 0$. By integrating with respect of

time t , $\dot{x} = C_1$. When $t=0$, $\dot{x} = U \cos 135^\circ$

$$= -\frac{U}{\sqrt{2}} \therefore C_1 = -\frac{U}{\sqrt{2}} \therefore \dot{x} = -\frac{U}{\sqrt{2}}$$

By integrating with respect of time t ,

$$x = -\frac{U}{\sqrt{2}}t + C_2 \text{ . When } t=0, x=2d$$

$$\therefore C_2 = 2d \therefore x = -\frac{U}{\sqrt{2}}t + 2d$$

Vertical motion of the projectile from A .

$\ddot{y} = -g$. By integrating with respect of

time t , $\dot{y} = -gt + K_1$. When $t=0$,

$$\dot{y} = U \sin 135^\circ = \frac{U}{\sqrt{2}} \therefore K_1 = \frac{U}{\sqrt{2}} \therefore \dot{y} = -gt + \frac{U}{\sqrt{2}}$$

By integrating with respect of time t .

$$y = -\frac{1}{2}gt^2 + \frac{U}{\sqrt{2}}t + K_2 \text{ . When } t=0, y=0$$

$$\therefore K_2 = 0 \therefore y = -\frac{1}{2}gt^2 + \frac{U}{\sqrt{2}}t \quad (3 \text{ marks})$$

iii) In order for the projectile to hit the

target, they have to have the same

coordinates at the same time. The equations

of the motions of the projectile from O

$$\text{are } x = \frac{V}{\sqrt{2}}t \text{ and } y = -\frac{1}{2}gt^2 + \frac{V}{\sqrt{2}}t$$

and the equations of the motions of the target are $x=d$ and

$$y = -\frac{1}{2}gt^2 + d$$

let $y=y$

$$\therefore -\frac{1}{2}gt^2 + \frac{V}{\sqrt{2}}t = -\frac{1}{2}gt^2 + d$$

$$t = \frac{\sqrt{2}d}{V} \text{ when } t = \frac{\sqrt{2}d}{V}$$

$x_{\text{target}} = d$ and $x_{\text{projectile from 0}}$

$$\text{is } x = \frac{V}{\sqrt{2}}t = \frac{V}{\sqrt{2}} \times \frac{\sqrt{2}d}{V} = d$$

$\therefore$ At $t = \frac{\sqrt{2}d}{V}$ the projectile

from 0 hits the target, as they

have the same co ordinates.

But in order for the impact to occur in the air it should be

$$y > 0 \text{ for } t = \frac{\sqrt{2}d}{V} \text{ But } y = -\frac{1}{2}gt^2 + d$$

$$\therefore -\frac{1}{2}g\left(\frac{\sqrt{2}d}{V}\right)^2 + d > 0$$

$$= \frac{1}{2} \times \frac{2d^2}{V^2} + d > 0 \quad -\frac{gd^2}{V^2} + d > 0$$

$$\text{(dividing by } d > 0) \quad -\frac{gd}{V^2} + 1 > 0$$

$$1 > \frac{gd}{V^2} \therefore V^2 > gd \therefore V > \sqrt{gd} \text{ (or } V > \sqrt{gd})$$

$\therefore$ The impact occurs in the air

$$V > \sqrt{gd} \quad (3 \text{ marks})$$

iv) After the impact, the fragment falls vertically down.

$\therefore$ The equations of the motion of the target can be used for the motion of the fragment. Now

the projectile from A hits the fragment from the same

co ordinate at the same time.

$$\text{For } y_{\text{projectile}} = y_{\text{fragment}}$$

$$-\frac{1}{2}gt^2 + \frac{U}{\sqrt{2}}t = -\frac{1}{2}gt^2 + d$$

$$= \frac{\sqrt{2}d}{U}$$

this time is the time taken for the projectile from A to hit the fragment.

The time taken for the projectile from 0 to hit the target is $t = \frac{\sqrt{2}d}{V}$

$$\therefore T = \text{time}_{\text{fragment}} - \text{time}_{\text{target}}$$

$$= \frac{\sqrt{2}d}{U} - \frac{\sqrt{2}d}{V}$$

$$= \sqrt{2}d \left(\frac{1}{U} - \frac{1}{V} \right) \quad (2 \text{ marks})$$