



Student Number	
Mark / 35	

Chemistry

Preliminary Course

Final Examination • 2005

General Instructions

- Reading time – 5 minutes
- Working time – 50 minutes
- Write using black or blue pen
- Draw diagrams using pencil
- Board-approved calculators may be used
- A Data Sheet and a Periodic Table are provided at the back of this paper and may be removed for convenience
- Write your Student Number at the top of this page

Total Marks – 35

Part A – 9 marks

- Attempt Questions 1 – 9
- Allow about 10 minutes for this part

Part B – 26 marks

- Attempt Questions 10 – 19
- Allow about 40 minutes for this part

Part A – 9 marks

Attempt Questions 1 – 9

Allow about 10 minutes for this part

Select the alternative A, B, C or D that best answers the question. Fill in the response oval completely.

Sample: $2 + 4 =$ (A) 2 (B) 6 (C) 8 (D) 9
A ☐ B ☒ C ☐ D ☐

If you think you have made a mistake, put a cross through the incorrect answer and fill in the new answer.

A ☒ B ☒ C ☐ D ☐

If you change your mind and have crossed out what you consider to be the correct answer, then indicate the correct answer by writing the word **correct** and drawing an arrow as follows.

A ☒ B ☒ C ☐ D ☐
correct

Answer Box for Questions 1 – 9

1	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
2	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
3	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
4	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
5	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
6	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
7	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
8	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>
9	A <input type="radio"/>	B <input type="radio"/>	C <input type="radio"/>	D <input type="radio"/>

1 Study the reaction... $\text{H}_2(\text{g}) + \text{Cl}_2(\text{g}) \rightarrow 2\text{HCl}(\text{g})$

Which of the statements cannot apply to this reaction?

- (A) One gram of H_2 reacts with one gram of Cl_2 to produce 2 grams of HCl .
- (B) One molecule of H_2 reacts with one molecule of Cl_2 to produce 2 molecules of HCl .
- (C) One litre of H_2 reacts with one litre of Cl_2 to produce 2 litres of HCl at constant conditions.
- (D) One mole of H_2 reacts with one mole of Cl_2 to produce 2 moles of HCl .

2 Which of the production sequences shows the extraction of copper from its ore to 99.9% pure copper?

- (A) froth flotation \rightarrow crushing and grinding \rightarrow smelting \rightarrow electrolysis
- (B) crushing and grinding \rightarrow froth flotation \rightarrow smelting \rightarrow electrolysis
- (C) crushing and grinding \rightarrow smelting \rightarrow froth flotation \rightarrow electrolysis
- (D) smelting \rightarrow crushing and grinding \rightarrow froth flotation \rightarrow electrolysis

3 Which of the following may **not** be a consequence of thermal pollution in water?

- (A) Reduction in dissolved oxygen.
- (B) Disruption of aquatic organisms breeding cycles.
- (C) Out of season migration of aquatic fauna.
- (D) Decrease in salt concentration.

4 At 100 kPa and 25°C , 4 litres of oxygen gas contain 1×10^{21} molecules.
Which of these gas samples also contains 1×10^{21} molecules under the same conditions?

- (A) 1 L of NH_3
- (B) 2 L of Cl_2
- (C) 4 L of CO_2
- (D) 8 L of He

5 To which reaction can Gay–Lussac’s law be applied?

- (A) $2\text{H}_2(\text{g}) + \text{O}_2(\text{g}) \rightarrow 2\text{H}_2\text{O}(\text{g})$
(B) $2\text{H}_2\text{O}(\text{l}) \rightarrow 2\text{H}_2(\text{g}) + \text{O}_2(\text{g})$
(C) $2\text{H}_2\text{O}(\text{l}) + 2\text{Na}(\text{s}) \rightarrow 2\text{Na}^+(\text{aq}) + 2\text{OH}^-(\text{aq}) + \text{H}_2(\text{g})$
(D) $\text{H}^+(\text{aq}) + \text{OH}^-(\text{aq}) \rightarrow \text{H}_2\text{O}(\text{l})$

6 Which property accounts for the moderate viscosity of water?

- (A) Specific heat capacity
(B) Hydrogen bonding
(C) Density
(D) Boiling point


7 Metals X, M, Z and D are all very useful metals. The table shows selected properties of the metals...

<i>Property</i>	<i>X</i>	<i>M</i>	<i>Z</i>	<i>D</i>
<i>Ionisation energy (kJ mol⁻¹)</i>	584	751	896	766
<i>Percentage Abundance in Earth’s Crust</i>	8	0.07	0.00001	0.07

Use the data to list the metals according to increasing market price.

- (A) $\text{X} < \text{M} < \text{D} < \text{Z}$
(B) $\text{D} < \text{X} < \text{Z} < \text{M}$
(C) $\text{X} < \text{Z} < \text{M} < \text{D}$
(D) $\text{Z} < \text{D} < \text{M} < \text{X}$

Answer Questions 8 and 9 using this reaction pathway diagram...



- 8 Which statement correctly describes this reaction?
- (A) The reaction is endothermic and the surroundings will become cooler.
 - (B) The reaction is exothermic and the surroundings will become cooler.
 - (C) The reaction is endothermic and the surroundings will become warmer.
 - (D) The reaction is exothermic and the surroundings will become warmer.
- 9 Which statement is true if a catalyst is added to the system?
- (A) ΔH remains constant and the activation energy increases.
 - (B) Both ΔH and activation energy decrease.
 - (C) ΔH remains constant and the activation energy decreases.
 - (D) Both ΔH and activation energy remain constant.

Part B – 26 marks


Attempt Questions 10 – 19

Allow about 40 minutes for this part

► *Show all relevant working in questions involving calculations.*

Question 10 (3 marks)

Air bags have saved thousands of lives and are now commonly fitted in new cars. An air bag inflates very rapidly by producing nitrogen gas from the decomposition of 100 grams of sodium azide...




- (a) Calculate the moles of sodium azide originally in the air bag. **(1 mark)**


- (b) Calculate the moles of nitrogen produced and the resultant volume at 100 kPa and 25°C. **(2 marks)**

Question 11 (3 marks)

Xenon tetrafluoride is an unstable compound which self-decomposes...



The graph shows the decomposition of a pure, 4 mole sample of XeF_4 over a period of 60 days.




- (a) Calculate the number of moles of XeF_4 which has decomposed after 16 days. **(1 mark)**

- (b) On which day are there equal moles of reactant and products present? **(1 mark)**

- (c) Calculate the mass of F_2 present at 40 days. **(1 mark)**

Question 12 (2 marks)

Iron will react with hydrochloric acid and appear to “dissolve” ...




Ken Chemiski places a 2.51 g iron nail in a beaker and prepares some dilute, 1.00 mol L^{-1} HCl.

Calculate the volume of acid required to fully react with the nail.

Question 13 (3 marks)

Héloïse prepares some beautiful golden lead(II) iodide crystals by this precipitation reaction...



She reacts 25.0 mL of 0.100 mol L^{-1} potassium iodide with excess lead(II) nitrate solution.

(a) Calculate the moles of lead(II) nitrate which reacted. **(2 marks)**

(b) Calculate the mass of PbI_2 produced. **(1 mark)**

Question 14 (3 marks)

In your studies, you performed a practical investigation to observe the effect of temperature on reaction rate. Describe your experiment, including the observed results.

Question 15 (2 marks)

- (a) Identify one pollutant produced by the incomplete combustion of an organic compound. **(1 mark)**

- (b) Write a chemical equation to show the incomplete combustion of butane (C_4H_{10}). **(1 mark)**

Question 16 (2 marks)

The diagram shows liquid methanol, CH_3OH , being poured into a beaker of water.



Use the symbols, $\text{H}-\text{O}-\text{H}$ for water and $\begin{array}{c} \text{H} & \text{H} \\ & \diagdown \quad \diagup \\ & \text{C} \\ & \diagup \quad \diagdown \\ \text{H} & \text{O}-\text{H} \end{array}$ for methanol to draw a diagram that illustrates the strongest intermolecular forces in the solution.

Question 17 (3 marks)

Outline the method required to prepare a 250 mL solution of 0.102 mol L⁻¹ strontium chloride.

► Write your answer in numbered sequential steps (1, 2, 3...).

us.com

Question 18 (2 marks)

Tincture of iodine, a common antiseptic, is a 2% (w/w) solution of iodine in ethanol.

- (a) Calculate the mass of iodine crystals is required to prepare a 250 g sample of tincture of iodine?
(1 mark)

- (b) Calculate the mass of ethanol is required? **(1 mark)**

Question 19 (3 marks)

In an experiment to measure the density of water at 28°C, an empty 100 mL measuring cylinder was found to have a mass of 150.5 grams. When 80.3 mL of water was poured into the cylinder, the mass of the cylinder and contents was 230.5 grams.

- (a) Calculate the density of water at 28°C based on the data. **(1 mark)**

- (b) When the measuring cylinder and water was sealed and kept inside a freezer overnight at -10°C , the volume reading increased to 85.9 mL. Explain this observation. **(1 mark)**

- (c) When water freezes it expands. Why is this process important in Nature? **(1 mark)**

DATA SHEET

Avogadro constant, N_A $6.022 \times 10^{23} \text{ mol}^{-1}$

Volume of 1 mole ideal gas: at 100 kPa and

at 0°C (273.15 K) 22.71 L

at 25°C (298.15 K) 24.79 L

Ionisation constant for water at 25°C (298.15 K), K_w 1.0×10^{-14}

Specific heat capacity of water $4.18 \times 10^3 \text{ J kg}^{-1} \text{ K}^{-1}$

Some useful formulae

$$\text{pH} = -\log_{10}[\text{H}^+]$$

$$\Delta H = -m C \Delta T$$

Some standard potentials

$\text{K}^+ + \text{e}^-$	\rightleftharpoons	K(s)	-2.94 V
$\text{Ba}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ba(s)	-2.91 V
$\text{Ca}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ca(s)	-2.87 V
$\text{Na}^+ + \text{e}^-$	\rightleftharpoons	Na(s)	-2.71 V
$\text{Mg}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mg(s)	-2.36 V
$\text{Al}^{3+} + 3\text{e}^-$	\rightleftharpoons	Al(s)	-1.68 V
$\text{Mn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Mn(s)	-1.18 V
$\text{H}_2\text{O} + \text{e}^-$	\rightleftharpoons	$\frac{1}{2}\text{H}_2(\text{g}) + \text{OH}^-$	-0.83 V
$\text{Zn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Zn(s)	-0.76 V
$\text{Fe}^{2+} + 2\text{e}^-$	\rightleftharpoons	Fe(s)	-0.44 V
$\text{Ni}^{2+} + 2\text{e}^-$	\rightleftharpoons	Ni(s)	-0.24 V
$\text{Sn}^{2+} + 2\text{e}^-$	\rightleftharpoons	Sn(s)	-0.14 V
$\text{Pb}^{2+} + 2\text{e}^-$	\rightleftharpoons	Pb(s)	-0.13 V
$\text{H}^+ + \text{e}^-$	\rightleftharpoons	$\frac{1}{2}\text{H}_2(\text{g})$	0.00 V
$\text{SO}_4^{2-} + 4\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	$\text{SO}_2(\text{aq}) + 2\text{H}_2\text{O}$	0.16 V
$\text{Cu}^{2+} + 2\text{e}^-$	\rightleftharpoons	Cu(s)	0.34 V
$\frac{1}{2}\text{O}_2(\text{g}) + \text{H}_2\text{O} + 2\text{e}^-$	\rightleftharpoons	2OH^-	0.40 V
$\text{Cu}^+ + \text{e}^-$	\rightleftharpoons	Cu(s)	0.52 V
$\frac{1}{2}\text{I}_2(\text{s}) + \text{e}^-$	\rightleftharpoons	I^-	0.54 V
$\frac{1}{2}\text{I}_2(\text{aq}) + \text{e}^-$	\rightleftharpoons	I^-	0.62 V
$\text{Fe}^{3+} + \text{e}^-$	\rightleftharpoons	Fe^{2+}	0.77 V
$\text{Ag}^+ + \text{e}^-$	\rightleftharpoons	Ag(s)	0.80 V
$\frac{1}{2}\text{Br}_2(\text{l}) + \text{e}^-$	\rightleftharpoons	Br^-	1.08 V
$\frac{1}{2}\text{Br}_2(\text{aq}) + \text{e}^-$	\rightleftharpoons	Br^-	1.10 V
$\frac{1}{2}\text{O}_2(\text{g}) + 2\text{H}^+ + 2\text{e}^-$	\rightleftharpoons	H_2O	1.23 V
$\frac{1}{2}\text{Cl}_2(\text{g}) + \text{e}^-$	\rightleftharpoons	Cl^-	1.36 V
$\frac{1}{2}\text{Cr}_2\text{O}_7^{2-} + 7\text{H}^+ + 3\text{e}^-$	\rightleftharpoons	$\text{Cr}^{3+} + \frac{7}{2}\text{H}_2\text{O}$	1.36 V
$\frac{1}{2}\text{Cl}_2(\text{aq}) + \text{e}^-$	\rightleftharpoons	Cl^-	1.40 V
$\text{MnO}_4^- + 8\text{H}^+ + 5\text{e}^-$	\rightleftharpoons	$\text{Mn}^{2+} + 4\text{H}_2\text{O}$	1.51 V
$\frac{1}{2}\text{F}_2(\text{g}) + \text{e}^-$	\rightleftharpoons	F^-	2.89 V

Aylward and Findlay, *SI Chemical Data* (5th Edition) is the principal source of data for this examination paper. Some data may have been modified for examination purposes.

KEY	
Atomic Number	79
Symbol of element	Au
Atomic Weight	197.0
Name of element	Gold

KEY	
Atomic Number	79
Symbol of element	Au
Atomic Weight	197.0
Name of element	Gold

57 La	58 Ce	59 Pr	60 Nd	61 Pm	62 Sm	63 Eu	64 Gd	65 Tb	66 Dy	67 Ho	68 Er	69 Tm	70 Yb	71 Lu
138.9	140.1	140.9	144.2	[144.9]	150.4	152.0	157.3	158.9	162.5	164.9	167.3	168.9	173.0	175.0
Lanthanum	Cerium	Praseodymium	Neodymium	Promethium	Samarium	Europium	Gadolinium	Terbium	Dysprosium	Holmium	Erbium	Thulium	Ytterbium	Lutetium

89 Ac [227.0]	90 Th 232.0	91 Pa 231.0	92 U 238.0	93 Np [237.0]	94 Pu [244.1]	95 Am [243.1]	96 Cm [247.1]	97 Bk [247.1]	98 Cf [251.1]	99 Es [252.1]	100 Fm [257.1]	101 Md [258.1]	102 No [259.1]	103 Lr [262.1]
Actinium	Thorium	Protactinium	Uranium	Neptunium	Plutonium	Americium	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lavrencium

Where the atomic weight is not known, the relative atomic mass of the most common radioactive isotope is shown in brackets. The atomic weights of Np and Tc are given for the isotopes ^{237}Np and ^{99}Tc .