

Independent.

2004

Higher School Certificate Trial Examination

Mathematics Extension 1

General Instructions

- Reading time – 5 minutes
- Working time – 2 hours
- Write using black or blue pen
- Board approved calculators may be used
- A table of standard integrals is provided with this paper
- All necessary working should be shown in every question

Total marks - 84

Attempt Questions 1 – 7


All questions are of equal value

This paper MUST NOT be removed from the examination room

Student name / number

Question 1	Begin a new page	Marks
------------	------------------	-------

a.	Find the domain of the function $f(x) = \log_e\left(\frac{5-x}{3-x}\right)$	3
b.	Consider the curves $y = x^3$ and $y = x^2 - x$.	
i.	Show that the curves intersect at the point where $x = 0$.	1
ii.	Find the acute angle between the curves at this point.	2
c.	If $P(x) = x^4 - 3x^3 + ax^2 - ax - 12$ is divisible by $(x - 3)$, find the value of a .	2
d.	In the diagram, $AB = 10$ cm, $CD = 15$ cm and $AC = x$. Find the value of x .	2



e.	Six people are to be placed around a circular table. Two of them want to sit together. In how many ways can the table be arranged?	2
----	---	---

Question 2	Begin a new page	Mark
a.	Find the exact value of $\int_{-1}^1 \sqrt{4 - x^2} dx$, using the substitution $x = 2\sin\theta$.	3
b.	For the expansion of the expression $\left(x - \frac{3}{x}\right)^5$, find the term independent of x	3
c.	i. Sketch the graph of $y = 2\tan^{-1} 3x$. ii. State the domain and range of the function.	2 1
d.	Solve the equation $3\cos\theta - 4\sin\theta = 5$, for $-\pi \leq \theta \leq \pi$. Express your answers to 2 decimal places.	3

Question 3	Begin a new page	
a.	In how many ways can 8 prefects be chosen from a group of 20 nominees?	1
b.	Find the exact value of $\int_0^{\pi/2} \sin^2 2x dx$.	3
c.	Use Mathematical Induction to show that $\sum_{r=1}^n 4r - 3 = 2n^2 - n$.	3
d.	Jaime wants to use Newton's Method to obtain the zero of $\sqrt[4]{x} = 0$. i. Using Newton's Method once with a first approximation of $x_1 = 1$, obtain a second approximation. ii. Using Newton's Method with a first approximation $x_1 \neq 0$, show that the second approximation, x_2 , is such that $ x_2 > x_1 $. Explain the significance of this result.	2 3

Question 4	Begin a new page	Marks
a.	A particle is moving so that its displacement, x cm, at any time, t seconds, is given by the equation $x = 2 \cos\left(3t + \frac{\pi}{6}\right)$.	
	i. Show that the particle moves in Simple Harmonic Motion.	2
	ii. State the period of the motion.	1
	iii. When does the particle first come to rest after $t = 0$?	1
b.	$P(2ap, ap^2)$ is a point on the parabola $x^2 = 4ay$. The normal at P cuts the x axis at S and the y axis at T .	
	i. Draw a half page sketch to show this information.	1
	ii. State the equation of the normal to the parabola at P and hence show that S is the point $(ap(2 + p^2), 0)$ and that T is the point $(0, a(2 + p^2))$.	3
	iii. Find the value(s) of p such that P is the midpoint of ST .	1
c.	i. Explain why the probability of obtaining 2 heads and a tail when three coins are tossed is $\frac{3}{8}$.	1
	ii. Sian tosses three coins 10 times in a row. Calculate the probability of obtaining 2 heads and a tail at least 2 times. Give your answer correct to 3 significant figures.	2

Question 5	Begin a new page	Marks
------------	------------------	-------

a. The rate at which a body cools in air is proportional to the difference between the temperature, T , of the body and the constant surrounding temperature, S . This can be expressed as


$$\frac{dT}{dt} = k(T - S)$$

where t is time in minutes and k is a constant.

i. Show that $T = S + Be^{-kt}$, where B is a constant, is a solution of the above equation. 1

ii. If a particular body cools from 100° to 80° in 30 minutes, find the temperature of the body after a further 30 minutes, given the surrounding temperature remains constant at 25° . Give your answer to the nearest degree. 3

b. A man of height 2 metres throws a ball from M to the roof of a 15 metre high building. He throws the ball at an initial velocity of 25m/s, and he is 20 m from the base of the building.



Between which two angles of projection must he throw the ball to ensure that it lands on the roof of the building? 5

(Assume $\ddot{x}=0$ and $\ddot{y}=-10$)

c. i. By considering $(1 + x)^{n+3} = (1 + x)^n(1 + x)^3$, show that 2

$$\binom{n+3}{k} = \binom{n}{k} + 3\binom{n}{k-1} + 3\binom{n}{k-2} + \binom{n}{k-3}.$$


ii. Between what values must k lie? 1

Question 6

Begin a new page

Marks

a. AB is a tangent to the circle. $AB \parallel EC$ and $CD = AC$.



Copy the diagram into your booklet.

Prove that $AC \parallel ED$.

b. A particle is moving in a straight line. At time t seconds, it has displacement x metres from a fixed point O on the line, velocity $v \text{ ms}^{-1}$ and acceleration $a \text{ ms}^{-2}$. The particle starts from O and you are given that $v = (2 - x)^2$.

i. Find an expression for a in terms of x . 1


ii. Find an expression for x in terms of t . 3

iii. Find the distance from O when the particle has a speed of 1 ms^{-1} . 1

c. i. Given a function, $y = f(x)$, under what geometrical conditions would $f(x) = f^{-1}(x)$? 2

ii. Give an example of a function for which $f(x) = f^{-1}(x)$. 1

Question 7	Begin a new page	Marks
a.	i. Find $\frac{d}{dx} \left(x \cos^{-1} x - \sqrt{1-x^2} \right)$	2
	ii. Find the area between the curve $y = \cos x$, the y axis and the lines $y = \frac{1}{2}$ and $y = \frac{\sqrt{3}}{2}$.	3
b.	The diagram shows a unit square, $ABCD$, where $A(1, 0)$, $B(1, 1)$, $C(2, 1)$, $D(2, 0)$.	



Copy the diagram into your workbook.

i. A line, l , passing through the origin with gradient m , cuts the sides AB and CD at P and Q respectively.

Comment on the possible values of m .

1

ii. For what value(s) of m does the line, l , divide the area of the square in the ratio 2:1?

3

iii. Another line, k , passes through the origin with gradient, n , and cuts the square through sides AB and BC at S and T respectively.

Show that it is not possible for k to divide the area of the square in the ratio 2:1.

3