



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2004**

**TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION**

# Mathematics Extension 1

## *General Instructions*

- Reading time – 5 minutes.
- Working time – 2 hours.
- Write using black or blue pen.
- Board approved calculators may be used.
- All necessary working should be shown in every question if full marks are to be awarded.
- Marks may **NOT** be awarded for messy or badly arranged work.
- Hand in your answer booklets in 3 bundles.  
Section A (Questions 1 - 3),  
Section B (Questions 4 - 5) and  
Section C (Questions 6 - 7).
- Start each Section in a **NEW** answer booklet.

## **Total Marks - 84 Marks**

- Attempt questions 1- 7
- All questions are of equal value.

Examiner: *R. Boros*

This is an assessment task only and does not necessarily reflect the content or format of the Higher School Certificate.

2

Total marks – 84

Attempt Questions 1 – 7

All questions are of equal value

Answer each section in a SEPARATE writing booklet. Extra writing booklets are available.

SECTION A (Use a SEPARATE writing booklet)

Question 1 (12 marks)

Marks

(a) Solve for  $x$ :  $(x^2 - 1)(x + 5) > 0$  2

(b) Differentiate  $y = \ln \sqrt{x+1}$  for  $x > -1$  2

(c) Use the Table of Integrals provided to evaluate 2

$$\int_0^{\frac{\pi}{6}} \sec 2x \tan 2x \, dx$$

(d) Find the exact value of  $\int_0^{\sqrt{3}} \frac{1}{9+x^2} \, dx$  2

(e) 8 people including A and B are to be seated around a circle. 2

How many arrangements are possible if A and B do not wish to sit together?


(f) Show that  $\frac{1 - \cos \theta}{\sin \theta} + \frac{\sin \theta}{1 + \cos \theta} = 2 \tan \frac{\theta}{2}$  2

Question 2 (12 marks)

Marks

- (a) Differentiate  $y = \sin^{-1} 2x$  2
- (b) Find the domain and range of  $y = 3 \sin^{-1} \sqrt{1-x^2}$  2
- (c) (i) Express  $\sqrt{3} \cos x - \sin x$  in the form  $R \cos(x + \alpha)$ ,  
where  $R > 0$  and  $0 < \alpha < \frac{\pi}{2}$ . 2
- (ii) Hence or otherwise, find the general solutions for  
 $\sqrt{3} \cos x - \sin x = 1$  2

- (d) In the diagram below  $ABCD$  is a cyclic quadrilateral.  
 $BA$  is produced to  $F$ .  
 $BC \parallel FE$   
 $CF$  and  $AE$  meet at  $D$ .



Copy or trace the diagram into your answer booklet.

- (i) Show that  $\triangle DEF \parallel \triangle FEA$  2
- (ii) Hence show that  $(EF)^2 = EA \times ED$  2

Section A is continued on page 4


## SECTION A continued

Question 3 (12 marks)

Marks

- (a) Use the Principle of Mathematical Induction to show that  $2^{3n} - 1$  is divisible by 7 for all integers  $n \geq 1$ . 3
- (b) For the curve  $y = 1 + 2 \cos x - 2 \cos^2 x$ ,
- (i) Show that  $\frac{dy}{dx} = 2 \sin x (2 \cos x - 1)$  1
- (ii) Hence find the stationary point(s) in the interval  $-\frac{\pi}{6} \leq x \leq \frac{\pi}{2}$  2
- (iii) Sketch the curve and find the greatest and least value of  $y$  in  $-\frac{\pi}{6} \leq x \leq \frac{\pi}{2}$  2

(c)



A rectangle is inscribed under the curve  $y = \frac{1}{1+x^2}$ , as shown in the diagram above, such that the rectangle is symmetrical about the  $y$  axis.

- (i) Show that the area of the rectangle is given by  $\frac{2a}{1+a^2}$ . 1
- (ii) Find the maximum area of the rectangle. 3

END OF SECTION A

(5)

**SECTION B (Use a SEPARATE writing booklet)**

**Question 4 (12 marks)**

**Marks**

- (a) (i) Show that the equation of the tangent at  $T(-2t, t^2)$  on the parabola  $y = \frac{1}{4}x^2$  is given by  $tx + y + t^2 = 0$ . 2
- (ii)  $M(x, y)$  is the midpoint of the interval  $TA$  where  $A$  is the  $x$  intercept of the tangent at  $T$ . 2
- Find the equation of the locus of  $M$  as  $T$  moves on the parabola.
- (b) Solve  $4x^3 - 12x^2 + 11x - 3 = 0$  if the roots are the terms of an arithmetic series. 3
- (c) (i) Find the point of intersection of the curves  $y = 2 \cos x$  and  $y = \frac{1}{2} \sec x$  in the interval  $-\frac{\pi}{2} < x < \frac{\pi}{2}$ . 2
- (ii) The area enclosed between the two curves listed above is rotated  $360^\circ$  about the  $x$  axis. 3
- Find the volume of the solid of revolution.  
(Leave your answer in exact form.)

**Section B is continued on page 6**

6

**SECTION B continued**

Question 5 (12 marks)

Marks

- (a) A spherical balloon leaks air such that the radius decreases at a rate of 5 cm/second. 2

Calculate the rate of change of the volume of the balloon when the radius is 100 mm.

[The volume of a sphere is  $V = \frac{4}{3}\pi r^3$ ]

- (b) A particle moves in such a way that its displacement  $x$  cm from the origin  $O$  after a time  $t$  seconds is given by

$$x = 2 \cos\left(t + \frac{\pi}{6}\right) \text{ cm}$$

- (i) Show that the particle moves in Simple Harmonic Motion. 2
- (ii) Evaluate the period of the motion. 1
- (iii) Find the time at which the particle first passes through the origin on its first oscillation. 1
- (iv) Find the velocity when the particle is 1 cm from the origin on its first oscillation. 2

- (c) Find  $\int \sqrt{16-x^2} dx$  using the substitution  $x = 4 \sin \theta$ . 4

**END OF SECTION B**

## SECTION C (Use a SEPARATE writing booklet)

Question 6 (12 marks)

Marks

- (a) Find a primitive function for  $\frac{3x}{4+x^2}$  1
- (b) If  $P(x) = 8x^3 - 12x^2 + 6x + 13$ ,
- (i) For what values of  $x$  is  $P(x)$  increasing? 1
- (ii) Show that  $P(x)$  has only one zero,  $x_1$  and that  $x_1 < 0$ . 1
- (iii) Taking  $x = -1$  as a first approximation to  $P(x) = 0$ , find a better approximation for  $x_1$ , using Newton's Method once. 2
- [Express your answer correct to 2 decimal places.]
- (c) At any time  $t$ , the rate of cooling of the temperature  $T$  of a body, when the surrounding temperature is  $S$ , is given by the differential equation
- $$\frac{dT}{dt} = -k(T - S)$$
- for some constant  $k$ .
- (i) Show that  $T = S + Ae^{-kt}$ , for some constant  $A$ , satisfies this differential equation. 2
- (ii) A metal rod has a temperature of  $1390^\circ \text{C}$  and cools to  $1060^\circ \text{C}$  in 10 minutes when the surrounding temperature is  $30^\circ \text{C}$ . 3
- Find how much *longer* it will take the rod to cool to  $110^\circ \text{C}$ , giving your answer to the nearest minute.
- (iii) Sketch the graph of the function  $T = S + Ae^{-kt}$ . 2

Section C continues on page 8



**SECTION C continued**

Question 7 (12 marks)

**Marks**

- (a) (i) A particle is projected from a point  $O$  with a velocity  $V$  at an angle  $\theta$  to the horizontal. 2

Taking the coordinate axes at the point of projection, find the parametric expressions for the velocity and the position of the particle at any time  $t$ .

[Take  $g = 10 \text{ m/s}^2$ ]

- (ii) After 1 second, the position of the particle is  $(6\sqrt{3}, 1)$ . 2

Show that the initial velocity and the angle of projection are 12 m/s and  $30^\circ$  respectively.

- (iii) Find the range of the motion. 2

- (b) In the expansion of  $\left(1 - \frac{2x}{3}\right)^7$ , state the coefficient of  $x^5$ . 2

- (c) If  $(1+x)^n = \sum_{k=0}^n {}^nC_k x^k$  find

(i)  $\sum_{k=1}^n {}^nC_k$  2

(ii)  $\sum_{k=1}^n k {}^nC_k$  2

**End of paper**



9

# STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, n \neq -1; x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax,$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, a > 0, -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE:  $\ln x = \log_e x, x > 0$



**SYDNEY BOYS HIGH SCHOOL**  
MOORE PARK, SURRY HILLS

**2004**

**TRIAL HIGHER SCHOOL  
CERTIFICATE EXAMINATION**

# Mathematics Extension 1

## Sample Solutions

Section	Marker
A	Mr Dunn
B	Ms Nesbitt
C	Mr Bigelow