



NORMANHURST BOYS' HIGH SCHOOL  
NEW SOUTH WALES

Student Number

--	--	--	--	--	--	--	--

Class: 12M 1 2 3 4 5 (Please Circle)

**2013**

**HIGHER SCHOOL CERTIFICATE  
TRIAL EXAMINATION**

# Mathematics Extension 1

## General Instructions

- Reading time - 5 minutes
- Working time - 2 hours
- Write using black or blue pen
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11-14, show relevant mathematical reasoning and/or calculations
- Begin each question on a separate writing booklet

## Total marks - 70

### Section I

Pages 2-5

### 10 marks

- Attempt Questions 1-10
- Answer on the Multiple Choice answer sheet provided
- Allow about 15 minutes for this section

### Section II

Pages 6-11

### 60 marks

- Attempt Questions 11-14
- Allow about 1 hour 45 minutes for this section

*Students are advised that this is a school-based examination only and cannot in any way guarantee the content or format of future Higher School Certificate Examinations.*

## Section I

10 marks

Attempt Questions 1-10

Allow about 15 minutes for this section

Use the multiple-choice answer sheet for Questions 1-10

---

- (1) The height of a giraffe has been modelled using the equation:

$$H = 5.40 - 4.80e^{-kt}$$

where  $H$  is the height in metres,  $t$  is the age in years and  $k$  is a positive constant.

If a 6 years old giraffe has a height of 5.16 metres, find the value of  $k$ , correct to 2 significant figures.

- (A) 0.05
- (B) 0.24
- (C) 0.50
- (D) 4.8

- (2) What is the value of  $\lim_{x \rightarrow 0} \left( \frac{\sin \frac{1}{3}x}{2x} \right)$

- (A)  $\frac{1}{6}$
- (B)  $\frac{2}{3}$
- (C)  $\frac{3}{2}$
- (D) 6


- (3) Which of the following equates to the expression  $\frac{1 - e^{3x}}{1 - e^{2x}}$ .

- (A)  $1 + \frac{e^{2x}}{1 + e^x}$
- (B)  $1 - e^x$
- (C)  $1 + e^x + e^{2x}$
- (D) None of the above


- (4) The point  $P$  divides the interval  $A(\frac{17}{3}, 2)$  to  $B(-3, 4)$  **externally** in the ratio  $2:3$ . Which one of the following is the coordinates of point  $P$ ?
- (A)  $(-23, 2)$
- (B)  $(-9, -12)$
- (C)  $(9, 0)$
- (D)  $(23, -2)$
- (5) A curve is defined by the parametric equations  $x = \sin 2t$  and  $y = \cos 2t$ . Which of the following, in terms of  $t$ , equates to  $\frac{dy}{dx}$ ?
- (A)  $-\tan 2t$
- (B)  $2 \tan 2t$
- (C)  $2 \sin 4t$
- (D)  $\cos 4t$
- (6) Which of the following is the inverse function of  $y = \frac{x-4}{x-2}$ ,  $x \neq 2$ ?
- (A)  $y = \frac{x-2}{x-4}$
- (B)  $y = f^{-1}(y)$
- (C)  $y = \frac{2(x-2)}{x-1}$
- (D)  $y = \frac{x+4}{x+2}$

(7) Which of the following represents the graph of  $y = \cos^{-1}(x+1)$ .


(A)




(B)



(C)



(D)



(8) Given that  $xy = x + 1$ , the definite integral  $\int_3^5 x \, dy$  equates to:

(A)  $e^2$

(B)  $\ln 2$

(C)  $-\ln\left(\frac{5}{3}\right)$

(D)  $e^{\frac{5}{3}}$

(9) The motion of a particle moving along the  $x$  – axis executes simple harmonic motion. The maximum velocity of the particle is  $4 \text{ m/s}$  and the period of motion is  $\pi$  seconds . Which of the following could be the displacement equation for this particle?

(A)  $x = 4 \cos \pi t$

(B)  $x = -\sin 2t$

(C)  $x = 2 \cos 2t$

(D)  $x = 2 + \cos 2t$

(10) A particle moves with a velocity  $v \text{ m/s}$  where  $v = \sqrt{x^2 + 1}$  . Given that  $x > 0$  , which of the following is equal to the acceleration of the particle when  $v = 4 \text{ m/s}$  .

(A)  $\sqrt{17} \text{ m/s}^2$

(B)  $-3 \text{ m/s}^2$

(C)  $\sqrt{15} \text{ m/s}^2$

(D)  $2\sqrt{17} \text{ m/s}^2$

## Section II

**60 marks**

**Attempt Questions 11-14**

**Allow about 1 hour and 45 minutes for this section**

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.

In Questions 11-14, your responses should include relevant mathematical reasoning and/or calculations.

---


**Question 11** (15 marks) Use a SEPARATE writing booklet.

- (a) Find the exact value of  $\sin 75^\circ$ . 2
- (b) Given that the acute angle between the lines  $y = mx$  and  $2x - 3y = 0$  is  $45^\circ$ , find possible value(s) of  $m$ . 3
- (c) Using the substitution  $u = 1 + x^2$ , or otherwise, evaluate 3
- $$\int_0^{\sqrt{8}} \left( \frac{x}{\sqrt{1+x^2}} \right) dx.$$
- (d) Solve the following inequality for  $x$ : 3
- $$\frac{1}{x} + \frac{x}{(x-2)} < 0$$
- (e) (i) On the same number plane, graph the following functions: 2
- $$y = 4 - x^2 \quad \text{and} \quad y = |3x|$$
- (ii) Hence or otherwise solve  $4 - x^2 \leq |3x|$  2

**Question 12** (15 marks) Use a SEPARATE writing booklet.

- (a) Find the exact value of  $\sin\left(2\cos^{-1}\frac{\sqrt{3}}{4}\right)$ . 2

- (b)  $AB$  is a chord of a circle centre  $O$ .  $AO$  is a diameter of a circle centre  $Q$ .  $Z$  is the point where the circle centre  $Q$  meets  $AB$ .




NOT  
SAE

- (i) Explain why  $AO = OB$ . 1
- (ii) Hence or otherwise, prove that  $AZ = ZB$ . 2
- (c) The quadratic equation  $x^2 - 4x + 9 = 0$  has roots  $\tan A$  and  $\tan B$ . Hence, find the size(s) of  $\angle(A + B)$ , noting that  $0 \leq A + B \leq 360^\circ$  (leave your answer to the nearest degree). 3
- (d) (i) By use of long division, find the remainder, in terms of  $a$  and  $b$  when  $P(x) = x^4 + 3x^3 + 6x^2 + ax + b$  is divided by  $x^2 + 2x + 1$ . 2
- (ii) If this remainder is  $3x + 2$ , find the values of  $a$  and  $b$ . 1
- (e) (i) Prove that  $\sin A \cos A \cos 2A = \frac{1}{4} \sin 4A$ . 2
- (ii) Hence or otherwise solve  $\sin A \cos A \cos 2A = 0$ , for  $0 \leq A \leq \frac{\pi}{2}$ . 2

**Question 13** (15 marks) Use a SEPARATE writing booklet.

- (a) Prove by mathematical induction that  $5^n \geq 1 + 4n$  for all integers  $n \geq 1$ . **3**

(b)



$P(2ap, ap^2)$  and  $Q(-2ap, ap^2)$  are variable points on the parabola  $x^2 = 4ay$ . The line  $PQ$  is parallel to the  $x$ -axis. The tangent at  $P$  meets the  $x$ -axis at  $Z$ .

- (i) Show that the equation of the tangent at  $P$  is given by  $y = px - ap^2$  **2**
- (ii) Hence show that  $Z = (ap, 0)$ . **1**
- (iii) Find the locus of midpoints of  $QZ$ . **2**
- (c) (i) Graph the function  $y = 2 \tan^{-1}(x)$ . **1**
- (ii) Graphically show why  $2 \tan^{-1}(x) - \frac{x}{4} = 0$  has one root, for  $x > 0$ . **1**
- (iii) Taking  $x_1 = 10$  as a first approximation to this root, use one application of Newton's method to find a better approximation, correct to 2 decimal places. **2**


**Question 13 continues on page 9**



Question 13 (continued)

13 (d)

3



A ladder AB, 5 metres long, is leaning against a vertical wall OA, with its foot B, on horizontal ground OB. The distances OB and OA are  $x$  and  $y$  metres respectively.  $x$  and  $y$  are related by the equation  $x^2 + y^2 = 25$ .

The foot of the ladder begins to slide along the ground away from the wall at a constant speed of 1 metre per second.

Find the speed at which the top of the ladder A is moving down the wall at the time when the top of the ladder is 4 metres above the ground.

**End of Question 13**


**Question 14** (15 marks) Use a SEPARATE writing booklet.

- (a) A particle is travelling in a straight line. Its displacement ( $x$  cm) from  $O$  at a given time ( $t$  sec) after the start of motion is given by:  $x = 2 + \sin^2 t$ .
- (i) Prove that the particle is undergoing simple harmonic motion. 2
- (ii) Find the centre of motion. 1
- (iii) Find the total distance travelled by the particle in the first  $\frac{3\pi}{2}$  seconds. 2
- (b) A shade sail with corners  $A$ ,  $B$  and  $C$  is shown in diagram 1, supported by three vertical posts. The posts at corners  $A$  and  $C$  are the same height, and the post at corner  $B$  is 2.4 m taller. Diagram 2 shows the sail in more detail.  $D$  is the point on the taller post horizontally level with the tops of the other two posts.  $AD = 6.4$  m and  $DC = 5.2$  m.  $\angle ADC = 125^\circ$ .  
Find the area of the shade sail  $ABC$  (leave your answer to 1 decimal place) 3

**Question 14 continues on page 11**

Question 14 (continued)

- (c) Over 80 years ago, during training exercises, the Army fired an experimental missile from the top of a building 15 m high with initial velocity ( $v$ ) where  $v = 130 \text{ m/s}$ , at an angle ( $\alpha$ ) to the horizontal. Noting that  $\alpha = \tan^{-1}\left(\frac{5}{12}\right)$  and taking  $g = 10 \text{ m/s}^2$



Assume that the equations of motion of the missile are  $\ddot{x} = 0$  and  $\ddot{y} = -10$

- (i) Show that  $\dot{x} = 120$  and  $\dot{y} = -10t + 50$ . 2  
Hence write down the equations of  $x$  and  $y$ .
- (ii) The rocket hit its intended target when its velocity reached  $60\sqrt{5} \text{ m/s}$ . 2  
Find the horizontal distance that the missile travelled to hit its target.
- (iii) The rocket was designed to hit its target once the angle to the horizontal of its flight path in a downward direction lies between  $20^\circ$  and  $30^\circ$ . Find the range of times after firing that this could happen. 3

**End of paper**

## STANDARD INTEGRALS

$$\int x^n dx = \frac{1}{n+1} x^{n+1}, \quad n \neq -1; \quad x \neq 0, \text{ if } n < 0$$

$$\int \frac{1}{x} dx = \ln x, \quad x > 0$$

$$\int e^{ax} dx = \frac{1}{a} e^{ax}, \quad a \neq 0$$

$$\int \cos ax dx = \frac{1}{a} \sin ax, \quad a \neq 0$$

$$\int \sin ax dx = -\frac{1}{a} \cos ax, \quad a \neq 0$$

$$\int \sec^2 ax dx = \frac{1}{a} \tan ax, \quad a \neq 0$$

$$\int \sec ax \tan ax dx = \frac{1}{a} \sec ax, \quad a \neq 0$$

$$\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a}, \quad a \neq 0$$

$$\int \frac{1}{\sqrt{a^2 - x^2}} dx = \sin^{-1} \frac{x}{a}, \quad a > 0, \quad -a < x < a$$

$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \ln \left( x + \sqrt{x^2 - a^2} \right), \quad x > a > 0$$

$$\int \frac{1}{\sqrt{x^2 + a^2}} dx = \ln \left( x + \sqrt{x^2 + a^2} \right)$$

NOTE:  $\ln x = \log_e x, \quad x > 0$